arXiv:1402.6074v1 [math.AT] 25 Feb 2014

THE SECONDARY PERIODIC ELEMENT β_{p^2/p^2-1} AND ITS APPLICATIONS

JIANGUO HONG AND XIANGJUN WANG

Abstract. In this paper, we prove that β_{p^2/p^2-1} survives to E_∞ in the Adams-Novikov spectral sequence $(ANSS)$ for all $p \geqslant 5$. As an easy consequence, we prove that $\beta_{sp^2/j}$ are permanent cycles for $s \geqslant 1$, $j \leqslant p^2-1$. From the Thom map $\Phi : Ext^{s,t}_{BP_*BP}(BP_*, BP_*) \longrightarrow Ext^{s,t}_{A}(\mathbb{Z}/p, \mathbb{Z}/p)$, we also see that h_0h_3 survives to E_∞ in the classical Adams spectral sequence.

1. Introduction

Let $p \geqslant 5$ be an odd prime. The Adams-Novikov spectral sequence $(ANSS)$ based on the Brown-Peterson spectrum is one of the most powerful tools to compute the p-component of the stable homotopy groups of spheres π_*S^0 (*cf.* [1, 4, 8, 20]). The E_2-term of the $ANSS$ is $Ext^{s,t}_{BP_*BP}(BP_*, BP_*)$.

From [10, 8] we know that $Ext^1_{BP_*BP}(BP_*, BP_*) = H^1BP_*$ is generated by $\alpha_{sp^n/n+1}$ for $n \geqslant 0$, $p \nmid s \geqslant 1$, where $\alpha_{sp^n/n+1}$ has order p^{n+1}. $Ext^2_{BP_*BP}(BP_*, BP_*) = H^2BP_*$ is the direct sum of cyclic groups generated by $\beta_{sp^n/j,i+1}$ for suitable (n, s, j, i) (*cf.* [8, 20, 21]).

It is known that each element $\alpha_{sp^n/n+1}$ in H^1BP_* is a permanent cycle in the $ANSS$ which represents an element of im$J$ having the same order. But we are far from fully determining which element of $\beta_{sp^n/j,i+1}$ in H^2BP_* survives to E_∞.

Let $\beta_{sp^n/j}$ denote $\beta_{sp^n/j,1}$. H. Toda [24, 25] proved $\alpha_1\beta_1^p$ is zero in π_*S^0. This relation supports a non-trivial Adams-Novikov differential $d_{2p-1}(\beta_{p/p}) = a\alpha_1\beta_1^p$, which is called the Toda differential. Based on Toda differential, D. Ravenel [17] proved that

$$d_{2p-1}(\beta_{p^n/p^n}) \equiv a\alpha_1\beta_{p^{n-1}/p^{n-1}} \mod \ker \beta_1^{p(p^{n-1}-1)/(p-1)}$$

for $n \geqslant 1$. That is to say, β_{p^n/p^n} can not survive to E_∞ in the Adams-Novikov spectral sequence. From this one can see that only $\beta_{sp^n/j} \in H^2BP_*$ for $s \geqslant 2$, $1 \leqslant j \leqslant p^n$ or $s = 1$, $1 \leqslant j \leqslant p^n - 1$ might survive to E_∞ in the $ANSS$. The following are some known results in this area:

Oka [11] proved that for $s = 1$, $1 \leqslant j \leqslant p-1$ or $s \geqslant 2$, $1 \leqslant j \leqslant p$, $\beta_{sp/j}$ is a permanent cycle in the $ANSS$.

Oka [13] proved that for $s = 1$, $1 \leqslant j \leqslant 2p-2$ or $s \geqslant 2$, $1 \leqslant j \leqslant 2p$, $\beta_{sp^2/j}$ is a permanent cycle in the $ANSS$.

Later Oka [15, 16] generalized the result to $n \geqslant 2$, i.e. for $n \geqslant 2$; $s = 1$, $1 \leqslant j \leqslant 2^{n-1}(p-1)$ or $s \geqslant 2$, $1 \leqslant j \leqslant 2^{n-1}p$, $\beta_{sp^n/j}$ survives to E_∞ in the $ANSS$.

Shimomura [23] proved that for $s \geqslant 1$, $1 \leqslant j \leqslant p^2-2$, $\beta_{sp^2/j}$ survives to E_∞ in the $ANSS$.

In this paper, we proved:

2000 *Mathematics Subject Classification.* Primary 55Q10, 55T15, 55Txx.

Key words and phrases. stable homotopy groups of spheres, Adams-Novikov spectral sequence, infinite descent method.

The authors were supported by NSFC grant No. 11071125, 11261062 and SRFDP.

Theorem A *Let $p \geqslant 5$ be an odd prime. Then β_{p^2/p^2-1} is a permanent cycle in the Adams-Novikov spectral sequence.*

Let M be the cofiber of the degree p map $p : S^0 \to S^0$,

$$S^0 \xrightarrow{p} S^0 \longrightarrow M.$$

There exists the Smith-Toda map $v_1^j : \Sigma^{|v_1^j|}M \to M$ and its cofiber is denoted by $M(1;j)$

$$\Sigma^{|v_1^j|}M \xrightarrow{v_1^j} M \longrightarrow M(1,j)\ .$$

D. Ravenel proved that

Theorem 7.12 [22] *Let $p \geqslant 5$ be an odd prime. If for some fixed $n \geqslant 1$,*

(i) *the spectrum $M(1, p^n - 1)$ is a ring spectrum,*
(ii) *β_{p^n/p^n-1} is a permanent cycle and*
(iii) *the corresponding homotopy element has order p,*

then $\beta_{sp^n/j}$ is a permanent cycle (and the corresponding homotopy element has order p) for all $s \geqslant 1$ and $1 \leqslant j \leqslant p^n - 1$.

From [14, 15, 16], we know that $M(1, p^n - 1)$ is a ring spectrum for $n \geqslant 1$, $p \geqslant 5$. Thus from the theorem above and Theorem A, we have:

Proposition B *Let $p \geqslant 5$ be an odd prime. Then for $s \geqslant 1$, $j \leqslant p^2-1$, $\beta_{sp^2/j}$ is a permanent cycle in the $ANSS$.*

It is not yet known that whether β_{sp^2/p^2} for $s \geqslant 2$ is a permanent cycle.

Between the $ANSS$ and the classical Adams spectral sequence (ASS), there is the Thom reduction map

$$\Phi : Ext^*_{BP_*BP}(BP_*, BP_*) \longrightarrow Ext^*_A(\mathbb{Z}/p, \mathbb{Z}/p)$$

and $\Phi(\beta_{p^2/p^2-1}) = h_0h_3$. Thus

Corollary C *Let $p \geqslant 5$ be an odd prime. Then h_0h_3 is a permanent cycle in the Adams spectral sequence.*

This paper is arranged as follows. In section 2 we recall the construction of the topological small descent spectral sequence ($TSDSS$) which is a spectral sequence that converges to π_*S^0 started from the homotogy groups of a complex with p-cells. Then we describe the generators of its E_1-term. In section 3 we compute a differential in the $TSDSS$, which is used in proving theorem A. Then in section 4, we prove our main theorem by the topological small descent spectral sequence.

2. The small descent spectral sequence and the ABC Theorem

In 1985, D. Ravenel [18, 19, 20, 21] introduced the method of infinite descent and used it to compute the first thousand stems of the stable homotopy groups of spheres at the prime 5. This method is an approach to finding the E_2-term of the $ANSS$ by the following spectral sequence referred as the small descent spectral sequence ($SDSS$).

Hereafter we set that $q = 2p - 2$. Let $T(n)$ be the Ranevel spectrum (*cf.* [20] Section 5, Chapter 6) characterized by

$$BP_*T(n) = BP_*[t_1, t_2, \cdots, t_n].$$

Then we have the following diagram

$$S^0 = T(0) \longrightarrow T(1) \longrightarrow T(2) \longrightarrow \cdots \longrightarrow T(n) \longrightarrow \cdots \longrightarrow BP,$$

where S^0 denote the sphere spectrum localized at an odd prime $p \geqslant 5$. Let $T(0)_{p-1}$ and $T(0)_{p-2}$ denote the $q(p-1)$ and $q(p-2)$ skeletons of $T(1)$ respectively, they are denoted by Y and $\overline{Y}$ for simple. Then

$$Y = S^0 \cup_{\alpha_1} e^q \cup \cdots \cup_{\alpha_1} e^{(p-2)q} \cup_{\alpha_1} e^{(p-1)q} \qquad \text{and} \qquad \overline{Y} = S^0 \cup_{\alpha_1} e^q \cup \cdots \cup_{\alpha_1} e^{(p-2)q}.$$

The BP-homologies of them are

$$BP_*(Y) = BP_*[t_1]/\langle t_1^p \rangle \qquad \text{and} \qquad BP_*(\overline{Y}) = BP_*[t_1]/\langle t_1^{p-1} \rangle.$$

From the definition above we get the following cofibre sequences

$$S^0 \xrightarrow{i'} Y \xrightarrow{j'} \Sigma^q \overline{Y} \xrightarrow{k'} S^1, \tag{2.1}$$

$$\overline{Y} \xrightarrow{i''} Y \xrightarrow{j''} S^{(p-1)q} \xrightarrow{k''} \Sigma \overline{Y}, \tag{2.2}$$

and the short exact sequences of BP_* homologies

$$0 \longrightarrow BP_* S^0 \xrightarrow{i'_*} BP_* Y \xrightarrow{j'_*} BP_* \Sigma^q \overline{Y} \longrightarrow 0, \tag{2.3}$$

$$0 \longrightarrow BP_* \overline{Y} \xrightarrow{i''_*} BP_* Y \xrightarrow{j''_*} BP_* S^{(p-1)q} \longrightarrow 0. \tag{2.4}$$

Put (2.3) and (2.4) together, one has the following long exact sequence

$$0 \longrightarrow BP_* S^0 \longrightarrow BP_*(Y) \longrightarrow BP_*(\Sigma^q Y) \longrightarrow BP_*(\Sigma^{pq} Y) \longrightarrow \cdots. \tag{2.5}$$

Put (2.1) and (2.2) together, one has the following Adams diagram of cofibres

$$\begin{array}{ccccccc} S^0 & \longleftarrow & \Sigma^{q-1}\overline{Y} & \longleftarrow & S^{pq-2} & \longleftarrow & \Sigma^{(p+1)q-3}\overline{Y} \longleftarrow \cdots \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ Y & & \Sigma^{q-1} Y & & \Sigma^{pq-2} Y & & \Sigma^{(p+1)q-3} Y. \end{array} \tag{2.6}$$

Then one has:

Proposition 2.1 *(Ravenel* [20] *7.4.2 Proposition) Let Y be as above. Then*

(a) *There is a spectral sequence converging to $Ext^{s+u,*}_{BP_*BP}(BP_*, BP_*(S^0))$ with E_1-term*

$$E_1^{s,t,u} = Ext^{s,t}_{BP_*BP}(BP_*, BP_*Y) \otimes E[\alpha_1] \otimes P[\beta_1], \quad \text{where}$$
$$\alpha_1 \in E_1^{0,q,1}, \qquad \beta_1 \in E_1^{0,qp,2}$$

and $d_r : E_r^{s,t,u} \longrightarrow E_r^{s-r+1,t,u+r}$. Where $E[-]$ denotes the exterior algebra and $P[-]$ denotes the polynomial algebra on the indicated generators. This spectral sequence is referred as the small descent spectral sequence (SDSS).

(b) *There is a spectral sequence converging to $\pi_*(S^0)$ with E_1-term*

$$E_1^{s,t} = \pi_*(Y) \otimes E[\alpha_1] \otimes P[\beta_1], \quad \text{where}$$
$$\alpha_1 \in E_1^{1,q}, \qquad \beta_1 \in E_1^{2,pq}$$

and $d_r : E_r^{s,t} \longrightarrow E_r^{s+r,t+r-1}$. This spectral sequence is referred as the topological small descent spectral sequence (TSDSS).

The above two spectral sequences produce the 0-line and the 1-line $Ext^{0,*}_{BP_*BP}(BP_*, BP_*(S^0))$, $Ext^{1,*}_{BP_*BP}(BP_*, BP_*(S^0))$ or the corresponding elements in $\pi_*(S^0)$ by $Ext^{0,*}_{BP_*BP}(BP_*, BP_*Y)$ and $Ext^{1,*}_{BP_*BP}(BP_*, BP_*Y)$. $Ext^{s,*}_{BP_*BP}(BP_*, BP_*(S^0))$ $(s \geqslant 2)$ or the corresponding elements in $\pi_*(S^0)$ is produced by $Ext^{s,*}_{BP_*BP}(BP_*, BP_*Y)$ $(s \geqslant 2)$ as described as the following ABC Theorem.

ABC Theorem *(7.4.3 ABC Theorem* [20]*) For $p > 2$ and $t - s < q(p^3 + p - 1) - 3$, $s \geqslant 2$*

$$Ext^{s,t}_{BP_*BP}(BP_*, BP_*Y) = A \oplus B \oplus C,$$

where A is the $\mathbb{Z}/p$-vector space spanned by

$$\begin{aligned} A =& \{\beta_{ip},\ \beta_{ip+1} | i \leqslant p-1\} \cup \left\{\beta_{p^2/p^2-j} | 0 \leqslant j \leqslant p-1\right\}, \\ B =& R \otimes \{\gamma_k | k \geqslant 2\} \end{aligned}$$

where

$$R = P[b_{20}^p] \otimes E[h_{20}] \otimes \mathbb{Z}/p \left\{\left\{b_{11}^k | 0 \leqslant k \leqslant p-1\right\} \cup \left\{h_{11} b_{20}^k | 0 \leqslant k \leqslant p-2\right\}\right\},$$

and

$$C^{s,t} = \bigoplus_{i \geqslant 0} R^{s+2i, t+i(p^2-1)q}.$$

From the generators of R, we can obtain the generators of C as follows:
Let $i = jp + m$. Then from $R^{s+2i,t+i(p^2-1)} \subset C^{s,t}$ we have

(1) $b_{20}^{(j+1)p} \in R^{2(p-m)+2(\underline{jp+m}), t+(\underline{jp+m})(p^2-1)q} \subset C^{2(p-m),t}$ is represented by

$$b_{20}^{p-m-1} u_{jp+m}$$

for $p - 1 \geqslant m \geqslant 1$. From which we have

$$b_{20}^{p-m-1} u_{jp+m} \otimes E[h_{20}] \otimes \left\{b_{11}^k | 0 \leqslant k \leqslant p-1\right\} \cup \left\{h_{11} b_{20}^k | 0 \leqslant k \leqslant p-2\right\},$$

where

$$u_{jp+m} \in C^{2, q\left[(j+1)p^2 + (j+m+1)p + m\right]}.$$

(2) $b_{11}^k b_{20}^{jp} \in R^{2(k-m)+2(\underline{jp+m}), t+(\underline{jp+m})(p^2-1)q} \subset C^{2(k-m),t}$ is represented by

$$b_{11}^{k-m-1} \beta_{(j+1)p/p-m}$$

for $p - 1 \geqslant k \geqslant m + 1 \geqslant 1$. From which we have

$$b_{11}^{k-m-1} \beta_{(j+1)p/p-m} \otimes E[h_{20}],$$

where

$$\beta_{(j+1)p/p-m} \in C^{2, q\left[(j+1)p^2 + jp + m\right]}.$$

- Especially $h_{20} b_{11}^{p-1} b_{20}^{jp} \in R^{3+2(\underline{jp+p-2}), t+(\underline{jp+p-2})(p^2-1)q} \subset C^{3,t}$ is represented by $h_{11} \beta_{(j+1)p/1,2}$, which is an element of order p^2.

(3) $h_{11} b_{20}^k b_{20}^{jp} \in R^{2(k-m)+1+2(\underline{jp+m}), t+(\underline{jp+m})(p^2-1)q} \subset C^{2(k-m)+1,t}$ is represented by

$$b_{20}^{k-m-1} \eta_{jp+m+1}$$

for $p - 2 \geqslant k \geqslant m + 1 \geqslant 1$, where

$$\eta_{jp+m+1} = h_{11} u_{jp+m} \in C^{3, q\left[(j+1)p^2 + (j+m+2)p + m\right]}.$$

(4) $h_{20}h_{11}b_{20}^k b_{20}^{jp} \in R^{2(k-m+1)+2(\underline{jp+m}),t+(\underline{jp+m})(p^2-1)q} \subset C^{2(k-m+1)t}$ is represented by

$$b_{20}^{k-m}\beta_{jp+m+2}$$

for $p-2 \geqslant k \geqslant m \geqslant 0$, where

$$\beta_{jp+m+2} \in C^{2,q\left[jp^2+(j+m+2)p+m+1\right]}.$$

- Especially $h_{20}h_{11}b_{20}^{p-2}b_{20}^{jp} \in R^{2+2(\underline{jp+p-2}),t+(\underline{jp+p-2})(p^2-1)q} \subset C^{2,t}$ is represented by $\beta_{(j+1)p/1,2}$, which is an element of order p^2.

From the ABC Theorem above, we compute that $Ext_{BP_*BP}^{s,t}(BP_*, BP_*Y)$ for $s \geqslant 2$, $t-s < q(p^3+p-1)-3$ is the $\mathbb{Z}_{(p)}$-module generated by the following generators, here the generators are listed as generators, total degree $t-s$ and $t-s$ *mod* $pq-2$, range of index; where $pq-2$ is the total degree of $\beta_1 \in E_1^{0,qp,2}$ in the $SDSS$.

Generators of A

Generators		$t-s$ and $t-s$ *mod* $pq-2$	Range of index
β_{ip}		$q[ip^2+ip-1]-2$	
	$\equiv$	$2(i-1)p+2i$	if $i \leqslant p-2$
	$\equiv$	0	if $i=p-1$
β_{ip+1}		$q[ip^2+(i+1)p]-2$	
	$\equiv$	$2ip+2i$	if $i \leqslant p-2$
	$\equiv$	$\underline{2p}_{2p}$	if $i=p-1$
β_{p^2/p^2-j}		$q[p^3+j]-2$	
	$\equiv$	$\underline{2(j+1)p-2j}_{2p}$	if $j \leqslant p-2$
	$\equiv$	4	if $j=p-1$

Generators of B

Generators	$t-s$ and $t-s$ mod $pq-2$	Range of index
$h_{11}b_{20}^k\gamma_i$	$q[(i+k)p^2+(i+k)p+i-2]-2k-4$	
	$\equiv 2(k+2i-2)p$	if $2 \leqslant i$, $k+2i \leqslant p$
	$\equiv 2(k+2i-p-1)p+2$	if $k+2i > p$
$h_{20}h_{11}b_{2,0}^k\gamma_i$	$q[(i+k)p^2+(i+k+1)p+i-1]-2k-5$	
	$\equiv 2(k+2i-1)p-1$	if $k+2i < p$
	$\equiv 2(k+2i-p)p+1$	if $k+2i \geqslant p$
$b_{11}^k\gamma_i$	$q[(i+k)p^2+(i-1)p+i-2]-2k-3$	
	$\equiv 2(k+2i-2)p-2k-1$	if $k+2i \leqslant p+1$
	$\equiv 1$	if $k=0$, $2i=p+1$
	$\equiv \underline{2(k+2i-p-1)p-2k+1}_{4p-3}$	if $k+2i \geqslant p+2$
$h_{20}b_{11}^k\gamma_i$	$q[(i+k)p^2+ip+i-1]-2k-4$	
	$\equiv 2(k+2i-1)p-2(k+1)$	if $k+2i \leqslant p$
	$\equiv \underline{2(k+2i-p)p-2k}_{2p}$	if $k+2i > p$

Generators of C

Generators	$t-s$ and $t-s$ mod $pq-2$	Range of index
$b_{11}^k b_{20}^{p-m-1} u_{jp+m}$	$q[(p-m+j+k+1)p^2+jp+m]-2(p-m+k)$	
	$\equiv 2(j+k+1)p+2(j-k+1)$	if $1 \leqslant m \leqslant p-1$
$h_{20} b_{11}^k b_{2,0}^{p-m-1} u_{jp+m}$	$q[(p-m+j+k+1)p^2+(j+1)p+m+1]-2(p-m+k)-1$	
	$\equiv 2(j+k+2)p+2(j-k+1)-1$	if $j+k \leqslant p-4$ or $j+k=p-3, 2j<p-5$
	$\equiv 2(j-k+2)p-1$	if $j+k=p-3, 2j \geqslant p-5$
$h_{11} b_{20}^{k+p-m-1} u_{jp+m}$	$q[(p-m+j+k+1)p^2+(j+k+1)p+m]-2(p-m+k)-1$	
	$\equiv 2(j+k+2)p+2(j-p)+3$	
$h_{20} h_{11} b_{2,0}^{k+p-m-1} u_{jp+m}$	$q[(p-m+j+k+1)p^2+(j+k+2)p+m+1]-2(p-m+k+1)$	
	$\equiv 2(j+k+2)p+2(j-p)+2$	
$b_{11}^{k-m-1}\beta_{(j+1)p/p-m}$	$q[(j+k-m)p^2+jp+m]-(2k-2m)$	
	$\equiv 2(j+k)p+2(j-k)$	if $j+k \leqslant p-2$ or $j+k=p-1$, $2j<p-1$
	$\underline{\equiv 2(j+k-p+1)p+2(j-k+1)}_{2p}$	if $j+k \geqslant p$ or $j+k=p-1$, $2j \geq p-1$
$h_{20} b_{11}^{k-m-1}\beta_{(j+1)p/p-m}$	$q[(j+k-m)p^2+(j+1)p+m+1]-(2k-2m+1)$	
	$\underline{\equiv 2(j+k+1)p+2(j-k)-1}_{4p-3}$	if $j+k \leqslant p-3$ or $j+k=p-2$, $2j \leqslant p-1$
	$\equiv 2(j+k-p+2)p+2(j-k)+1$	if $j+k>p-2$ or $j+k=p-2$, $2j>p-1$
$h_{1,1}\beta_{(j+1)p/1,2}$	$q[(j+1)p^2+(j+2)p-1]-3$	
	$\equiv 2jp+2(j+1)+1$	if $j \leqslant p-3$
	$\equiv 1$	if $j=p-2$
$b_{2,0}^{k-m-1}\eta_{jp+m+1}$	$q[(j+k-m)p^2+(j+k+1)p+m]-(2k-2m+1)$	
	$\equiv 2(j+k)p+2j+1$	if $j+k \leqslant p-2$
	$\underline{\equiv 2(j+k-p+2)p+2(j-p)+3}_{4p-3}$	if $2p-5 \geqslant j+k>p-2$
$b_{2,0}^{k-m}\beta_{jp+m+2}$	$q[(j+k-m)p^2+(j+k+2)p+m+1]-2(k-m+1)$	
	$\underline{\equiv 2(j+k+1)p+2j}_{2p}$	if $j+k \leqslant p-3$
	$\equiv 2(j+k-p+3)p+2(j-p)+2$	if $j+k>p-3$
	$\equiv 0$	if $j=p-2$, $k=p-3$
$\beta_{(j+1)p/1,2}$	$q[(j+1)p^2+(j+1)p-1]-2$	
	$\equiv 2jp+2(j+1)$	if $j \leqslant p-3$
	$\equiv 0$	if $j=p-2$

Remark. The Adams-Novikov spectral sequence for the spectrum Y collapses from E_2-term $Ext^{s,t}_{BP_*BP}(BP_*, BP_*Y)$ in the range $t-s < q(p^3+p-1)-3$, since there are no elements with filtration $> 2p$. Thus we actually get the homotopy groups $\pi_{t-s}(Y)$ in this range.

3. A differential in the $TSDSS$

This section is armed at showing that

$$d_{2p-1}(h_{20}b_{11}\gamma_s) = \alpha_1\beta_1^p h_{20}\gamma_s$$

in the $TSDSS$, which will be used in proving Theorem A in section 4.

We begin from showing that $\pi_{q(p^2+2p+2)-2}V(2) = 0$. From which we show that the Toda bracket $\langle \alpha_1\beta_1, p, \gamma_s\rangle = 0$ and the Toda bracket $\langle \alpha_1\beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s\rangle$ is well defined. Then from that relation

$$\langle \alpha_1\beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s\rangle = \alpha_1\beta_1^{p-1}h_{20}\gamma_s = \beta_{p/p-1}\gamma_s.$$

and $d(h_{20}b_{11}) = \beta_1\beta_{p/p-1}$ we get the desired differential in the $TSDSS$.

Let $p \geqslant 5$ be an odd prime and $V(2)$ be the Smith-Toda spectrum characterized by

$$BP_*V(2) = BP_*/I_3$$

where I_3 is the invariant ideal of $BP_* = \mathbb{Z}_{(p)}[v_1, v_2, \cdots, v_i, \cdots]$ generated by $\langle p, v_1, v_2\rangle$. To compute the homotopy groups of $V(2)$, one has the $ANSS$ $\{E_r^{s,t}V(2), d_r\}$ that converges to $\pi_*V(2)$. The E_2-term of this spectral sequence is

$$E_2^{s,t}V(2) = Ext^{s,t}_{BP_*BP}(BP_*, BP_*M)$$

Let (BP_*, Γ) be a Hopf algebroid and M be a Γ-comodule with coaction $\psi : M \to M \otimes_{BP_*} \Gamma$, one has cobar complex of M given by

$$C^s_\Gamma M = M \otimes_{BP_*} \Gamma \otimes_{BP_*} \Gamma \otimes_{BP_*} \cdots \otimes_{BP_*} \Gamma$$

(with s factors of Γ). The differential $d : C^s_\Gamma M \to C^{s+1}_\Gamma M$ is of degree $+1$ given by

$$\begin{aligned} d(m \otimes x_1 \otimes \cdots \otimes x_s) &= \psi(m) \otimes x_1 \otimes \cdots \otimes x_s \\ &\quad + \sum_{i=1}^{s} (-1)^i m \otimes x_1 \otimes \cdots \otimes x_{i-1} \otimes \Delta(x_i) \otimes x_{i+1} \otimes \cdots \otimes x_s, \end{aligned} \tag{3.1}$$

where $\Delta : \Gamma \longrightarrow \Gamma \otimes_{BP_*} \Gamma$ is the coproduct of Γ. The cohomology of $(C^s_\Gamma M, d)$ is $Ext^{s,*}_\Gamma(BP_*, M)$. Thus we have the cobar complex

$$C^s_{BP_*BP}BP_*/I_3 = BP_*/I_3 \otimes_{BP_*} BP_*BP \otimes_{BP_*} \cdots \otimes_{BP_*} BP_*BP$$

whose cohomology is $Ext^{s,*}_{BP_*BP}(BP_*, BP_*V(2))$.

Let

$$\Gamma = BP_*/I_3 \otimes_{BP_*} BP_*BP \otimes_{BP_*} BP_*/I_3 = BP_*/I_3[t_1, t_2, \cdots].$$

Then (BP_*, Γ) is a Hopf algebroid. The structure maps of (BP_*, Γ) is given by

$$\begin{aligned} \eta_R(v_3) &= v_3; \\ \Delta(t_1) &= t_1 \otimes 1 + 1 \otimes t_1, \\ \Delta(t_2) &= t_2 \otimes 1 + t_1 \otimes t_1^p + 1 \otimes t_2, \\ \Delta(t_3) &= t_3 \otimes 1 + t_2 \otimes t_1^{p^2} + t_1 \otimes t_2^p + 1 \otimes t_3. \end{aligned} \tag{3.2}$$

One can easily see that

$$\begin{aligned} C^s_{BP_*BP}BP_*V(2) =& BP_*/I_3 \otimes_{BP_*} BP_*BP \otimes_{BP_*} \cdots \otimes_{BP_*} BP_*BP \\ =& BP_*/I_3 \otimes_{BP_*} \Gamma \otimes_{BP_*} \cdots \otimes_{BP_*} \Gamma \\ =& C^s_\Gamma BP_*/I_3, \end{aligned}$$

and then

$$Ext^{s,t}_{BP_*BP}(BP_*, BP_*V(2)) = Ext^{s,t}_\Gamma(BP_*, BP_*/I_3)$$

Lemma 3.1. *The $q(p^2+2p+2)-2$ dimensional stable homology group of $V(2)$ is trivial, i.e.,*

$$\pi_{q(p^2+2p+2)-2}V(2) = 0.$$

Proof. Fix $t-s = q(p^2+2p+2)-2$, we know that the Adams-Novikov E_2-term

$$Ext^{s,s+q(p^2+2p+2)-2}_{BP_*BP}(BP_*, BP_*V(2)) = Ext^{s,s+q(p^2+2p+2)-2}_\Gamma(BP_*, BP_*/I_3)$$

converges to $\pi_{q(p^2+2p+2)-2}V(2)$. We will prove that $\pi_{q(p^2+2p+2)-2}V(2) = 0$ by showing that $Ext^{s,s+q(p^2+2p+2)-2}_{BP_*BP}(BP_*, BP_*V(2)) = 0$.

In the cobar complex $C^s_\Gamma BP_*/I_3$, the inner degree of $|v_i| = |t_i| \geqslant q(p^3+p^2+p+1)$ for $i \geqslant 4$. It follows that in the range $t-s \leqslant q(p^3+p^2+p+1)-1$,

$$Ext^{s,t}_{BP_*BP}(BP_*, BP_*/I_3) = Ext^{s,t}_\Gamma(BP_*, BP_*/I_3) = Ext^{s,t}_{\Gamma'}(BP_*, BP_*/I_3).$$

where $\Gamma' = \mathbb{Z}/p[v_3][t_1,t_2,t_3]$. From $\eta_R(v_3) = v_3$ in (3.2), we see that

$$Ext^{s,*}_{\mathbb{Z}/p[v_3][t_1,t_2,t_3]}(BP_*, BP_*/I_3) \cong Ext^{s,*}_{\mathbb{Z}/p[t_1,t_2,t_3]}(\mathbb{Z}/p, \mathbb{Z}/p) \otimes \mathbb{Z}/p[v_3].$$

To compute the Ext groups $Ext^*_{\mathbb{Z}/p[t_1,t_2,t_3]}(\mathbb{Z}/p, \mathbb{Z}/p)$, we can use the modified May spectral sequence (MSS) introduced in [6, 7, 21].

There is the May spectral sequence $\{E_r^{s,t,*}, \delta_r\}$ that converges to $Ext^{s,t}_{\mathbb{Z}/p[t_1,t_2,t_3]}(\mathbb{Z}/p, \mathbb{Z}/p)$. The E_1-term of this spectral sequence is

$$E_1^{*,*,*} = E[h_{ij} | 0 \leqslant j, i = 1,2,3,] \otimes P[b_{ij} | 0 \leqslant j, i = 1,2,3], \tag{3.3}$$

where

$$h_{ij} \in E_1^{1,q(1+p+\cdots+p^{i-1})p^j, 2i-1} \qquad \text{and} \qquad b_{ij} \in E_1^{2,q(1+p+\cdots+p^{i-1})p^{j+1}, p(2i-1)}.$$

The first May differential is given by

$$\delta_1(h_{i,j}) = \sum_{0<k<i} h_{i-k,k+j}h_{k,j} \quad \text{and} \quad \delta_1(b_{i,j}) = 0. \tag{3.4}$$

For the reason of the total degree, to compute $Ext^{s,s+(q(p^2+2p+2)-2)}_{BP_*BP}(BP_*, BP_*/I_3)$ we only need to consider the sub-module generated by $h_{30}, h_{20}, h_{10}, h_{21}, h_{11}, h_{12}$ and b_{20}, b_{10}, b_{11}, i. e. the subcomplex

$$E[h_{ij} | 1 \leqslant i, i+j \leqslant 3] \otimes E[b_{20}, b_{11}] \otimes P[b_{10}].$$

From (3.4), we know that the May's E_2-term

$$E_2^{s,*,*} = H^{s,*,*}(E_1^{s,*,*}, \delta_1) = H^{*,*,*}(E[h_{ij} | 0 \leqslant j, i+j \leqslant 3], \delta_1) \otimes E[b_{20}, b_{11}] \otimes P[b_{10}]$$

within $t-s \leqslant q(p^2+2p+2)-1$. H. Toda in [26] computed the cohomology of $(E[h_{ij} | 0 \leqslant j, i+j \leqslant 3], \delta_1)$ which is the $\mathbb{Z}/p$-module generated by the following elements:

dim 0	1				
dim 1	h_{10}	h_{11}	h_{12}		
dim 2	$h_{10}h_{12}$	$h_{20}h_{10}$	$h_{20}h_{11}$	$h_{21}h_{11}$	$h_{21}h_{12}$

dim 3 $h_{20}h_{20}h_{11}$ $h_{21}h_{11}h_{12}$ $h_{21}h_{20}h_{11}$ $h_{30}h_{12}h_{10}$ $h_{30}h_{20}h_{10}$ $h_{30}h_{21}h_{12}$
dim 4 $h_{30}h_{20}h_{10}h_{11}$ $h_{30}h_{20}h_{10}h_{12}$ $h_{30}h_{21}h_{12}h_{10}$ $h_{30}h_{21}h_{12}h_{11}$ $h_{30}h_{21}h_{20}h_{11}$
dim 5 $h_{30}h_{21}h_{20}h_{11}h_{10}$ $h_{30}h_{21}h_{20}h_{12}h_{10}$ $h_{30}h_{21}h_{20}h_{12}h_{11}$
dim 6 $h_{30}h_{21}h_{20}h_{12}h_{11}h_{10}$

Note that $q(p^2+2p+2)-2 \equiv 0 \ mod\ 2$, we list all the even dimensional generators and total degree $t-s$ of $H^*(E[h_{ij}|i+j \leqslant 3]) \otimes E[b_{20}, b_{11}]$ in the range $q(p^2+2p+2)-2$.

$h_{20}h_{10}$	$q(p+2)-2$;	$h_{20}h_{11}$	$q(2p+1)-2$;
$h_{10}h_{12}$	$q(p^2+1)-2$;	$h_{21}h_{11}$	$q(p^2+2p)-2$;
b_{20}	$q(p^2+p)-2$;	b_{11}	qp^2-2;
$h_{20}h_{10}b_{11}$	$q(p^2+p+2)-4$;	$h_{20}h_{11}b_{11}$	$q(p^2+2p_1)-4$;
$h_{20}h_{10}b_{20}$	$q(p^2+2p+2)-4$.		

Suppose we have a generator y in $Ext^{s,s+q(p^2+2p+2)-2}_{\mathbb{Z}/p[v_3][t_1,t_2,t_3]}(BP_*, BP_*/I_3)$. Then y is the form of x or v_3x where x is an even dimensional generator in $H^*(E[h_{ij}|i+j \leqslant 3]) \otimes E[b_{20}, b_{11}] \otimes P[b_{10}]$. From the total degree $t-s$ of b_{10} is $qp-2$, one can easily see that except for $h_{20}h_{10}$ and $h_{20}h_{11}$, for any generator x listed above the total degree $t-s$ of xb_{10}^k and $v_3xb_{10}^k$ is not $q(p^2+2p+2)-2$. Suppose we have a generator $h_{20}h_{10}b_{10}^k, h_{20}h_{11}b_{10}^k$ or b_{10}^k whose total degree is $q(p^2+2p+2)-2$, then the total degree of $h_{20}h_{10}, h_{20}h_{11}$ or 1 should be $q(p^2+2p+2)-2$ module the total of b_{10}. From

$$\begin{aligned} q(p^2+2p+2)-2 \equiv & 6p-2 \quad mod \quad qp-2 \\ q(p+2)-2 \equiv & 4p-4 \quad mod \quad qp-2 \\ q(2p+1)-2 \equiv & 2p \quad mod \quad qp-2 \end{aligned}$$

we know that the total degree of any generator of the form $h_{20}h_{10}b_{10}^k, h_{20}h_{11}b_{10}^k$ or b_{10}^k is not $q(p^2+2p+2)-2$. Thus

$$Ext^{s,s+q(p^2+2p+2)-2}_{BP_*BP}(BP_*, BP_*V(2)) = Ext^{s,s+q(p^2+2p+2)-2}_{\mathbb{Z}/p[v_3][t_1,t_2,t_3]}(BP_*, BP_*/I_3) = 0.$$

The Lemma follows. □

It is easily showed that the following theorem holds from the lemma above.

Theorem 3.2. *For $p \geqslant 7$, $s \geqslant 1$, the Toda bracket $\langle \alpha_1\beta_1, p, \gamma_s \rangle = 0$*

Proof. Let $\widetilde{v}_3$ be the composition of the following maps

$$S^{q(p^2+p+1)} \xrightarrow{\widetilde{i}} \Sigma^{q(p^2+p+1)}V(2) \xrightarrow{v_3} V(2)$$

where the first map is inclusion to the bottom cell.

It is known that $\widetilde{v}_3$ is an order p elements in $\pi_{q(p^2+p+1)}V(2)$. Thus the Toda bracket $\langle \alpha_1\beta_1, p, \widetilde{v}_3 \rangle$ is well defined and $\langle \alpha_1\beta_1, p, \widetilde{v}_3 \rangle \in \pi_{q(p^2+2p+2)-2}V(2) = 0$. It follows that the Toda bracket $\langle \alpha_1\beta_1, p, \widetilde{v}_3 \rangle = 0$

Let $\widetilde{j} : V(2) \longrightarrow S^{q(p+2)+3}$ be the collapsing lower cells map from $V(2)$, then $\gamma_s = \widetilde{v}_3 \cdot v_3^{s-1} \cdot \widetilde{j}$. As a result,

$$\langle \alpha_1\beta_1, p, \gamma_s \rangle = \langle \alpha_1\beta_1, p, \widetilde{v}_3 \cdot v_3^{s-1} \cdot \widetilde{j} \rangle = \langle \alpha_1\beta_1, p, \widetilde{v}_3 \rangle \cdot v_3^{s-1} \cdot \widetilde{j} = 0$$

because $\langle \alpha_1\beta_1, p, \widetilde{v}_3 \rangle = 0 \in \pi_{q(p^2+2p+2)-2}V(2) = 0$. □

Proposition 3.3. *(cf. [20] 7.5.11 and [21] 7.6.11) Let $p \geqslant 7$ be an odd prime. Then in π_*S^0, the Toda bracket $\langle \alpha_1\beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s \rangle$ is well defined and*

$$\alpha_1\beta_1^{p-1}h_{20}\gamma_s = \langle \alpha_1\beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s \rangle = \beta_{p/p-1}\gamma_s.$$

Proof. From $\langle \beta_1^{p-1}, \alpha_1\beta_1, p, \rangle = 0$, $\langle \alpha_1\beta_1, p, \alpha_1\rangle = 0$ $\langle \alpha_1, \alpha_1\beta_1, p\rangle\rangle = 0$ and $\langle \alpha_1\beta_1, p, \gamma_s\rangle = 0$, we know that the following four fold Toda bracket is well defined and

$$\beta_{p/p-1} = \langle \beta_1^{p-1}, \alpha_1\beta_1, p, \alpha_1\rangle; \qquad \alpha_1 h_{20}\gamma_s = \langle \alpha_1, \alpha_1\beta_1, p, \gamma_s\rangle.$$

On the other hand, one has

$$\begin{aligned}
\beta_1^{p-1}\alpha_1 h_{20}\gamma_s &= \beta_1^{p-1}\langle \alpha_1, \alpha_1\beta_1, p, \gamma_s\rangle \\
&= \langle \alpha_1\beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s\rangle \\
&= \alpha_1\langle \beta_1^{p-1}, \alpha_1\beta_1, p, \gamma_s\rangle \\
&= \langle \beta_1^{p-1}, \alpha_1\beta_1, p, \alpha_1\gamma_s\rangle \\
&= \langle \beta_1^{p-1}, \alpha_1\beta_1, p, \alpha_1\rangle \cdot \gamma_s \\
&= \beta_{p/p-1}\gamma_s
\end{aligned}$$

The the proposition holds. □

Theorem 3.4. *Let $p \geqslant 7$ be an odd prime and $2 \leqslant s \leqslant p-2$. Then in the topological small descent spectral sequence ($TSDSS$), we have the following differential*

$$d_{2p-1}(h_{2,0}b_{1,1}\gamma_s) = \alpha_1\beta_1^p h_{2,0}\gamma_s.$$

Proof. Recall from [21] page ?, one has the differential in the small descent spectral sequence ($SDSS$)

$$d_1(h_{20}b_{11}) = \beta_1\beta_{p/p-1}.$$

Then from $\beta_1\beta_{p/p-1}\gamma_s \in Ext^{7,*}_{BP_*BP}(BP_*, BP_*Y)$ and the relation $\alpha_1\beta_1^{p-1}h_{20}\gamma_s = \beta_{p/p-1}\gamma_s$, we the differential in the $TSDSS$

$$d_{2p-1}(h_{2,0}b_{1,1}\gamma_s) = \beta_1\cdot\beta_{p/p-1}\gamma_s = \beta_1\cdot\beta_1^{p-1}\alpha_1 h_{20}\gamma_s = \alpha_1\beta_1^p h_{20}\gamma_s.$$

The theorem follows. □

4. The proof of Theorem A

In this section we prove our main theorem by showing that β_{p^2/p^2-1} survives to E_∞ in the topological small descent spectral sequence.

Consider the $TSDSS$ introduced in section 2 Proposition 2.1 (b), the element $\beta_{p^2/p^2-1} \in E_1^{0,q(p^3+1)-2}$. To prove that β_{p^2/p^2-1} survives to E_∞ in the $TSDSS$, it is sufficient to prove that all the differentials $d_r(\beta_{p^2/p^2-1}) = 0$ for $r \geqslant 1$. From $\beta_{p^2/p^2-1} \in E_1^{0,q(p^3+1)-2}$, we see that

$$d_r(\beta_{p^2/p^2-1}) \in E_r^{r,q(p^3+1)-3+r},$$

in the $TSDSS$.

Lemma 4.1. *For $r \geqslant 1$, the E_1-term $E_1^{r,q(p^3+1)-3+r}$ of the $TSDSS$ is a $\mathbb{Z}/p$ module generated by the following 8 elements*

$$\begin{aligned}
&\mathfrak{g}_1 = \beta_1^{\frac{p^2-6p+1}{2}} b_{11}^2\gamma_{\frac{p+1}{2}} \in E_1^{p^2-6p+1,*}; && \mathfrak{g}_2 = \beta_1^{p^2-p}h_{20}\beta_{p/p} \in E_1^{2p^2-2p,*};\\
&\mathfrak{g}_3 = \beta_1^{p-1}\eta_{(p-3)p+3} \in E_1^{2p-2,*};\\
&\mathfrak{g}_4 = \alpha_1\beta_{(p-1)p+1} \in E_1^{1,q(p^3+1)-2}; && \mathfrak{g}_5 = \alpha_1\beta_{p^2/p^2} \in E_1^{1,q(p^3+1)-2};\\
&\mathfrak{g}_6 = \alpha_1\beta_1^{\frac{p^2-2p-1}{2}} h_{2,0}\gamma_{\frac{p+1}{2}} \in E_1^{p^2-2p,*}; && \mathfrak{g}_7 = \alpha_1\beta_1^{mp-\frac{p-1}{2}} b_{11}^{\frac{p-1}{2}-m}\beta_{(\frac{p+1}{2})p/p-m} \in E_1^{(2m-1)p+2,*};\\
&\mathfrak{g}_8 = \alpha_1\beta_1^{p^2-1}\beta_2 \in E_1^{2p^2-1,*}.
\end{aligned}$$

Proof. From the ABC Theorem, we know that the generators of $E_1^{r,q(p^3+1)-3+r}$ are of the form $W=\beta_1^k w$ or $W=\alpha_1\beta_1^k w$, where w is an element listed in the ABC Theorem.

1. If a generator of $E_1^{r,q(p^3+1)-3+r}$ is of the form $W=\beta_1^k w$, then the total degree of $\beta_1 w$ is $q(p^3+1)-3$ and the total degree of w is $q(p^3+1)-3$ module the total degree of β_1 $(qp-2)$. From $q(p^3+1)-3\equiv 4p-3$ *mod* $qp-2$, we list all the generators whose total degree is $4p-3$ *mod* $qp-2$ (marked with underline and subscript $4p-3$ in the ABC Theorem).

$$b_{11}^2\gamma_{\frac{p+1}{2}}, \qquad h_{20}\beta_{p/p}, \qquad \eta_{(p-3)p+3}.$$

From which we get the following generators in $E_1^{r,q(p^3+1)-3+r}$:

$$\mathfrak{g}_1=\beta_1^{\frac{p^2-6p+1}{2}}b_{11}^2\gamma_{\frac{p+1}{2}}; \qquad \mathfrak{g}_2=\beta_1^{p^2-p}h_{20}\beta_{p/p}; \qquad \mathfrak{g}_3=\beta_1^{p-1}\eta_{(p-3)p+3}.$$

2. If a generator of $E_1^{r,q(p^3+1)-3+r}$ is of the form $W=\alpha_1\beta_1^k w_1$, then the total degree of $\alpha_1 w_1$ must be congruent to $4p-3$ modulo $|\beta_1|=qp-2$, and the total degree of w_1 is $2p$ modulo $qp-2$ from $|\alpha_1|=2p-3$. We can find all such w_1's are as follows (marked with underline and subscript $2p$ in the ABC Theorem):

$$\beta_{(p-1)p+1}; \qquad \beta_{p^2/p^2}; \qquad h_{20}\gamma_{\frac{p+1}{2}}; \qquad b_{11}^{\frac{p-1}{2}-m}\beta_{(\frac{p+1}{2})p/p-m}; \qquad \beta_2.$$

From which we get the following generators in $E_1^{r,q(p^3+1)-3+r}$:

$$\mathfrak{g}_4=\alpha_1\beta_{(p-1)p+1}; \qquad \mathfrak{g}_5=\alpha_1\beta_{p^2/p^2}; \qquad \mathfrak{g}_6=\alpha_1\beta_1^{\frac{p^2-2p-1}{2}}h_{2,0}\gamma_{\frac{p+1}{2}};$$
$$\mathfrak{g}_7=\alpha_1\beta_1^{mp-\frac{p-1}{2}}b_{11}^{\frac{p-1}{2}-m}\beta_{(\frac{p+1}{2})p/p-m}; \qquad \mathfrak{g}_8=\alpha_1\beta_1^{p^2-1}\beta_2.$$

Compute the filtration of the corresponding generators, we get the lemma. □

Theorem 4.2. *In the topological small descent spectral sequence, the element*

$$\beta_{p^2/p^2-1}\in E_1^{0,q(p^3+1)-2}$$

is a permanent cycle.

This means that β_{p^2/p^2-1} is a permanent cycle in the ANSS by the relation between the TSDSS and the ANSS.

Proof. We know that $\beta_{p^2/p^2-1}\in Ext_{BP_*BP}^{2,*}(BP_*,BP_*)$ and in the $TSDSS$ $d_1(\beta_{p^2/p^2-1})=0$. Thus $\mathfrak{g}_4$ and $\mathfrak{g}_5$ have too low filtration to to be the target of $d_r\beta_{p^2/p^2-1}$.

In the $TSDSS$ we have the following differentials.

(1) From $d_{2p-1}(b_{11})=\alpha_1\beta_1^p$, we have

$$d_{2p-1}(\mathfrak{g}_1)=d_{2p-1}(\beta_1^{\frac{p^2-6p+1}{2}}b_{11}^2\gamma_{\frac{p+1}{2}})=2\alpha_1\beta_1^{\frac{p^2-4p+1}{2}}b_{11}\gamma_{\frac{p+1}{2}}.$$

(2) From $d_1(h_{20}\beta_{p/p})=\beta_1\beta_{p/p-1}$ (*cf.* [21] section 5 of Chapter 7), we have

$$d_1(\mathfrak{g}_2)=d_1(\beta_1^{p^2-p}h_{20}\beta_{p/p})=\beta_1^{p^2-p+1}\beta_{p/p-1}.$$

(3) From the Toda differential $d_{2p-1}(\beta_{p/p})=\alpha_1\beta_1^p$ and the relation $\alpha_1\eta_{(p-3)p+3}=\beta_{(p-3)p+4}\beta_{p/p}$ ([21] 7.5.7), we have

$$\begin{aligned}&d_{2p-1}(\alpha_1\eta_{(p-3)p+3})=\alpha_1\ d_{2p-1}\eta_{(p-3)p+3}\\ =&d_{2p-1}(\beta_{(p-3)p+4}\beta_{p/p})=\beta_{(p-3)p+4}\ d_{2p-1}(\beta_{p/p})=\alpha_1\beta_{(p-3)p+4}\beta_1^p.\end{aligned}$$

Thus $d_{2p-1}(\eta_{(p-3)p+3}) = \beta_{(p-3)p+4}\beta_1^p$ and

$$d_{2p-1}(\mathfrak{g}_3) = d_{2p-1}(\beta_1^{p-1}\eta_{(p-3)p+3}) = \beta_1^{2p-1}\beta_{(p-3)p+4}.$$

(4) From Theorem 3.4, we have

$$d_{2p-1}(\beta_1^{\frac{p^2-4p-1}{2}}h_{20}b_{11}\gamma_{\frac{p+1}{2}}) = \beta_1^{\frac{p^2-4p-1}{2}}\alpha_1\beta_1^p h_{20}\gamma_{\frac{p+1}{2}} = \alpha_1\beta_1^{\frac{p^2-2p-1}{2}}h_{20}\gamma_{\frac{p+1}{2}} = \mathfrak{g}_6.$$

(5) For $1 \leqslant m \leqslant \frac{p-1}{2}$, from $d_1(h_{20}\beta_{(\frac{p+1}{2})p/p-m+1}) = \beta_1\beta_{(\frac{p+1}{2})p/p-m}$ (*cf.* [21] section 5 of Chapter 7), we have

$$d_1(\mathfrak{g}_7) = d_1(\alpha_1\beta_1^{mp-1-\frac{p-1}{2}}h_{20}b_{11}^{\frac{p-1}{2}-m}\beta_{(\frac{p+1}{2})p/p-m+1}) = \alpha_1\beta_1^{mp-\frac{p-1}{2}}b_{11}^{\frac{p-1}{2}-m}\beta_{(\frac{p+1}{2})p/p-m}.$$

(6) We have

$$d_{2p-1}(\beta_1^{p^2-p-1}b_{11}\beta_2) = \alpha_1\beta_1^{p^2-1}\beta_2 = \mathfrak{g}_8$$



From the discussion above, we known that all the differentials $d_r(\beta_{p^2/p^2-1}) = 0$, for $r \geqslant 1$. This show that β_{p^2/p^2-1} is a permanent cycle in the $TSDSS$. □

Proposition 4.3. *Let $p \geqslant 5$ be an odd prime. For $s \geqslant 1$, $j \leqslant p^2-1$, $\beta_{sp^2/j}$ is a permanent cycle in the $ANSS$ and the corresponding homotopy element has order p.*

Proof. This proposition is easily got from the theorem above and the [22] Theorem 7.12. □

Corollary 4.4. *Let $p \geqslant 5$ be an odd prime. h_0h_3 is a permanent cycle in the Adams spectral sequence.*

Proof. Let $\phi : BP \to K\mathbb{Z}/p$ be the Thom map which induces the Thom reduction map

$$\Phi : Ext^*_{BP_*BP}(BP_*, BP_*) \longrightarrow Ext^*_A(\mathbb{Z}/p, \mathbb{Z}/p),$$

then $\Phi(\beta_{p^2/p^2-1} + x) = h_0h_3$, $x \in \ker\Phi$ (*cf.* [8] Theorem 9.4).

Hence h_0h_3 is the permanent cycle and the corresponding homotopy class has order p since $x = 0$ by degree reason. □

Remark At prime 5, D. Ravenel in [20] page 304 proved that $\beta^5_{5/5} = \beta_1 x_{952} \in \pi_{950}S^0$ survives to E_∞ in the $ANSS$. Then from $\langle \beta^5_{5/5}, \alpha_1, 5\rangle = 0$ and $\langle \alpha_1, 5, \alpha_1\rangle = 0$, we know that the four fold Toda bracket

$$\langle \beta^5_{5/5}, \alpha_1, 5, \alpha_1\rangle$$

is well defined and then $\beta_{5^2/5^2-1} = \langle \beta^5_{5/5}, \alpha_1, 5, \alpha_1\rangle$ survives to E_∞ in the $ANSS$.

References

[1] Adams, J. F., *On the strucure and applications of the Steenrod algebra*, Comm. Math. Helv.**32** (1958), 180-214.

[2] Cohen R., *Odd primary infinite families in stable homotopy theory*, Mem. Amer. Math. Soc. **30** (1981) no. 242 VIII +92pp.

[3] Kato, R., Shimomura, K., *Products of greek letter elements dug up from the third morava stabilizer algebra*, Algebr. Geom. Topol. **12** (2012), 951-961.

[4] Liulevicius, A., *The factorization of cyclic reduced powers by secondary cohomology operations*, Mem. Amer. Math. Soc. **42** (1962).

[5] Liu, X., Wang, X., *A four-filtered May spectral sequence and its applications,* Acta Math. Sin., (Engl. Ser.) **24** (2008), 1507-1524.

[6] May, J.P.,: *The cohomology of restricted Lie algebras and of Hopfalgebras; Applications to the Steenrod algebra (Theses)*, Princeton (1964).

[7] May, J.P.,: *The cohomology of restricted Lie algebras and of Hopf algebras*, J. Algebra **3** (1966), 123-146.

[8] Miller,H., Ravenel, D. C., Wilson, S., *Periodic phenomena in the Adams-Novikov spectral sequence*, Ann. of Math. **106** (1977), 469-516.

[9] Nakai, H., *An algebraic generalization of Image J* , Homology, Homotopy and Apllications. **10(3)** (2008), 321-333.

[10] Novikov, S. P., *The metods of algebraic topology from the viewpoint of cobordism theories*, Izv. Akad. Nauk. SSSR. Ser. Mat. **31** (1967), 855-951 (Russian).

[11] Oka, S., *A new family in the stable homotopy groups of sphere I*, Hiroshima Math. J. **5** (1975), 87-114.

[12] Oka, S., *A new family in the stable homotopy groups of sphere II*, Hiroshima Math. J. **6** (1976), 331-342.

[13] Oka, S., *Realizing some cyclic BP_*-modules and applications to stable homotopy of spheres*, Hiroshima Math. J. 7 (1977), 427-447.

[14] Oka, S., *Ring spectra with few cells*, Japan. J. Math. **5** (1979), 81-100.

[15] Oka, S., *Multiplicative structure of finite ring spectra and stable homotopy of spheres*, Algebraic Topology (Aarhus 1982) 41841. Lect. Notes in Math. **1051** Springer-Verlag 1984.

[16] Oka, S., *Small ring spectra and p-rank of the stable homotopy of spheres*, Contemp. Math. **19** (1983), 267-308.

[17] Ravenel, D. C., *The nonexistence of odd primary Arf invariant elements in stable homotopy theory*, Math. Proc. Cambridge Phil. Soc. *83* (1978), 429-443.

[18] Ravenel, D. C., The Adams-Novikov E_2-term for a complex with p-cells, Amer. J. Math. **107(4)** (1978), 933-968.

[19] Ravenel, D. C., *The method of infinite descent in stable homotopy theory. I* Recent progress in homotopy theory (Baltimore, MD, 2000), Contemp. Math., vol. 293, Amer. Math. Soc., Providence, RI, 2002, pp. 251-284.

[20] Ravenel, D. C., *Complex Cobordism and Stable Homotopy Groups of Spheres*, Academic Press, New York, 1986.

[21] Ravenel, D. C., *Complex cobordism and stable homotopy groups of spheres*, A. M. S. Chelsea Publishing, Providence, 2004.

[22] Ravenel, D. C., *A novice's guide to the Adams-Novikov spectral sequence*, Proc. Evanston Homotopy Theory Conf. Lect. Notes in Math., **658**, 404-475.

[23] Shimomura, K., *The beta elements $\beta_{tp^2/r}$ in the homotopy of spheres*, Algebr. Geom. Topol. **10** (2010) 2079-2090.

[24] Toda, H., *An important relation in the homotopy groups of spheres*, Proc. Japan Acad. **43** (1967), 893-942.

[25] Toda, H., *Extended p-th powers of complexes and applications to homotopy theory*, Proc. Japan Acad., **44** (1968), 198-203.

[26] H. Toda, *On spectra realizing exterior parts of Steenord algebra*, Topology **10** (1971), 55-65.

[27] Wang, X., *The secondary differentials on the third line of the Adams spectral sequence*, Topology. Appl. **156** (2009) 477-499.

Jianguo Hong, School of Mathematical Science and LPMC, Nankai University, Tianjin 300071, P. R. China
E-mail address: jghong66@163.com

Xiangjun Wang, School of Mathematical Science and LPMC, Nankai University, Tianjin 300071, P. R. China
E-mail address: xjwang@nankai.edu.cn